

實證醫學之開課須知及教學技巧介紹

如何將EBM運用於實際臨床教學-EBM 診斷型及治療型問題

衛生福利部雙和醫院
影像醫學部

蘇逸欣

2017.02.18

Outline

- 什麼是EBM?
- 流行病學簡介
- EBM五步驟

什麼是EBM?

- 請問什麼是EBM的縮寫?
- (1) Ebay, baby boss, MM chocolate
 - (2) Evidence-based medicine
 - (3) Emome, bb TV, Momo cartoons
 - (4) Esophagus burn medicine

什麼是EBM?

- EBM :

實證醫學 (EBM, Evidence-based medicine) 是以**流行病學**和**統計學**的方法，從龐大的醫學資料庫中嚴格評讀、綜合分析找出**值得信賴**的部分，並將所能獲得的**最佳文獻證據**，應用於**臨床工作**中，使病人獲得最佳的照顧。

Fieldm & Lohrm研究報告發現醫師每天所作的決策：

- 4%是有強而有力的臨床研究證據所支持
- 45%是有謹慎的臨床研究證據，且醫師間有一定程度的共識
- 51%不但缺乏有力的證據支持，在醫師間亦無共識，屬灰色地帶

什麼是EBM?

- EBM :
西元1972年，英國臨床流行病學者Archie Cochrane提出「謹慎地、明確地、小心地採用目前最佳的證據，作為照顧病人臨床決策的參考」。
- [Conscientious, explicit, and judicious use of current best evidence in making decisions about individual patients. ~ Archie Cochrane 1972]

什麼是EBM?

- EBM :
□ 實證醫學” 此一名詞自1992年加拿大McMaster大學Gordon Guyatt教授所領導的小組正式命名後，1992英國國家衛生部成立實證醫學中心，以Archie Cochrane之名命名，並由David L. Sackett擔任實證醫學中心主任，進而促成1993年Cochrane Collaboration的設立。



Dave Sackett

什麼是EBM?

● EBM :

- 目前世界各地有13個國家、15個實證醫學中心〔Cochrane Center〕在積極推動這個工作，其目標是從龐大的醫學資料庫中嚴格過濾、評讀這些文獻，並做系統性文獻回顧及綜合分析，以方便應用於臨床工作中，作為照護病患的依據。

什麼是EBM?

● EBM :



什麼是EBM?

● EBM :

56歲男性病人突然急性腹痛及背痛至急診就醫，做了KUB攝影發現左腎隱約有結石，因為影像不清楚因此又開立了IVP檢查，病人至影像醫學科檢查時問了問題：我從報導中得知低劑量電腦斷層也可以偵查出泌尿道結石，請問低劑量電腦斷層與IVP檢查有什麼差異以及他們的準確度為何？

什麼是EBM?

● 請問你要用什麼關鍵字來搜尋？

- (1) 食尚玩家, 浩角翔起
- (2) 腎結石, KUB
- (3) 腎結石, IVP, 低劑量電腦斷層
- (4) Running man, 金泰映

什麼是EBM?

● EBM :

Renal colic: A prospective evaluation of non-enhanced spiral CT versus intravenous pyelography.

Conclusions:

Non-enhanced spiral CT provided **greater diagnostic utility** in this randomized comparison **but no difference in measured outcomes.**

Mendelson, R. M. (2003) et al. Journal of Medical Image and Radiation Oncology

Outline

- 什麼是EBM?
- 流行病學簡介
- EBM五步驟

流行病學簡介

- **流行病學** (epidemiology) 是研究特定人群中**疾病**、健康狀況的分布及其決定因素，並研究防治疾病及促進健康的策略和措施的科學。搜尋證據 (Evidence search)
- 流行病學的定義和特徵。描述分布的常用測量指標 (**發病率**、**罹患率**、**患病率**、**死亡率**、**病死率**) 的意義、用途與計算方法。描述疾病**流行強度**的常用語：**爆發**、**流行**和**大流行**；疾病的分布的概念及主要內容。

什麼是EBM?

- 請問霍亂病毒主要感染途徑為？
 - (1) 眼神接觸以及四目交接
 - (2) 主要由飲水所傳染，污染來自嘔吐物或糞便，食物傳染但較少見
 - (3) 血液
 - (4) 接吻

流行病學簡介

John Snow與霍亂的大事記



- 流行病學發展的中心人物當屬**John Snow** (1813-1858) 這個人。他是一位臨床麻醉科醫師，但也是早期少數清楚瞭解到一個醫生的工作應該要超乎治療疾病的人之一。
- 透過一系列的觀察與研究，Snow發現1854年倫敦地區的**霍亂**流行是由**飲用水**受到**人類糞便**污染而引發的。

流行病學簡介

John Snow與霍亂的大事記

- 這項發現比微生物學家發現霍亂弧菌 (*Vibrio cholerae*) 的時間**早32年**；也比Pasteur於1865年證明**微生物**會造成**疾病流行**的觀察**早了10年**。
- 1854年英倫敦地區爆發霍亂流行，當時**倫敦**的飲水是由**數家私人水公司**經營。

流行病學簡介

John Snow與霍亂的大事記



- Snow分析了所有霍亂死亡個案的飲水來源後發現飲用**Southwark & Vauxhall**這家公司供水的人有較高的**死亡率**。
- 當數據顯示Southwark & Vauxhall公司所供應的家戶其霍亂死亡率是其他公司的**5至10倍**後，他開始懷疑霍亂流行是與Southwark & Vauxhall公司所供應的水有關。

流行病學簡介

John Snow與霍亂的大事記

- Snow正確的使用「**比率**」描述疾病在族群中分佈的情形是他能夠始終在正確方向上的重要因素。他接著仔細觀察Southwark & Vauxhall公司供水的來源，它發現**Southwark & Vauxhall**公司的水源位於**河川**的**下游**，河川上游並有**人類糞便**的排泄。

流行病學簡介

John Snow與霍亂的大事記

- 當Snow有了這些觀察後，他接續進行了幾個流行病學研究過程中很重要的步驟。
- 當他將霍亂病例按居住地區繪於地圖上時，他發現個案有「聚集」於Broadway Street上取水站的現象，而這也更令他相信霍亂的發生與水有關。
- 透過人時地的描述嘗試發現聚集 (clustering)現象是描述性流行病學研究的核心工作。

流行病學簡介

John Snow與霍亂的大事記

- Snow的這項觀察實際上是融入了現代科學研究中對照組 (control group)的概念，也使用了現代流行病學研究中經常使用的假說演繹 (hypothetico-deductive reasoning) 的推論。

流行病學簡介

John Snow與霍亂的大事記

- 雖然當時的生物學知識與技術並無法提供進一步的驗證支持Snow的發現，但當倫敦衛生當局於1854年9月8日將Broadway Street取水站的壓水手把移除後，霍亂死亡個案數隨即急速減少，到9月20日以後每日報告的死亡數便與9月1日暴發流行之前一樣。

流行病學簡介

John Snow與霍亂的大事記



流行病學簡介

近代流行病學

- 社區飲水加氯的試驗提供了能夠預防齲齒的氟濃度。
- 沙克疫苗的臨床試驗證據提供了人類消滅小兒麻痺的利器。
- Framingham Heart Study讓人類更加瞭解心血管疾病的病因與預防之道。
- 吸菸與肺癌及其他健康效應的研究。
- 長崎廣島原子彈爆炸存活者的追縱研究都讓人類更加瞭解環境與健康的互動。

Outline

- 什麼是EBM?
- 流行病學簡介
- EBM五步驟

什麼是EBM?

EBM五步驟：

- 提出問題 (Question formulation)
- 搜尋證據 (Evidence search)
- 嚴格判讀 (Critical appraisal)
- 恰當運用 (Evidence application)
- 評估結果 (Outcome evaluation)

EBM五步驟

EBM五步驟：

- 提出問題 (Question formulation)
- 搜尋證據 (Evidence search)
- 嚴格判讀 (Critical appraisal)
- 恰當運用 (Evidence application)
- 評估結果 (Outcome evaluation)

● 提出問題 (Question formulation)

How Questions Works

Find, Focus, Search

EBM五步驟

EBM五步驟：

- 提出問題 (Question formulation)
- 搜尋證據 (Evidence search)
- 嚴格判讀 (Critical appraisal)
- 恰當運用 (Evidence application)
- 評估結果 (Outcome evaluation)

● 提出問題 (Question formulation)

Background, Foreground

Novice → Expert

EBM五步驟

EBM五步驟：

- 提出問題 (Question formulation)
- 搜尋證據 (Evidence search)
- 嚴格判讀 (Critical appraisal)
- 恰當運用 (Evidence application)
- 評估結果 (Outcome evaluation)

● 提出問題 (Question formulation)

背景問題 (Background questions) : 5W1H
 意指與疾病相關的知識性問題，問題的結構通常包括 (who、what、where、when、how、why)。

Who : 病患特性、危險因子等
What : 自然病程、症狀學等
Where : 影響的器官與系統、疾病分佈的區域等
When : 疾病好發的年齡、時間、季節等
How : 病理生理學
Why : 病因

EBM五步驟

EBM五步驟：

- 提出問題 (Question formulation)
- 搜尋證據 (Evidence search)
- 嚴格判讀 (Critical appraisal)
- 恰當運用 (Evidence application)
- 評估結果 (Outcome evaluation)

● 提出問題 (Question formulation)

背景問題 (Background questions) :
 案例:急性心肌梗塞

Who : 老年人、男性、家族史
What : 胸痛、心電圖的變化等等
Where : 冠狀動脈阻塞的位置、程度等等
When : 氣溫低、心肌耗氧量高之活動等
How : 動脈硬化斑塊破裂、血栓形成
Why : 動脈硬化危險因子、糖尿病、高血壓、高血脂症、抽菸、肥胖, 少運動

EBM五步驟

EBM五步驟：

- 提出問題 (Question formulation)
- 搜尋證據 (Evidence search)
- 嚴格判讀 (Critical appraisal)
- 恰當運用 (Evidence application)
- 評估結果 (Outcome evaluation)

● 提出問題 (Question formulation)

前景問題 (Foreground questions) :
 指與治療病患相關的特殊性問題，通常包括有 **6 個** 項目：

Therapy/Prevention : 評估不同治療方式...
Harm/Etiology : 評估會造成的傷害...
Diagnosis (tests) : 不同診斷工具的準確度...
Prognosis : 評估疾病癒後...

EBM五步驟

EBM五步驟：
 提出問題 (Question formulation)
 搜尋證據 (Evidence search)
 嚴格判讀 (Critical appraisal)
 恰當運用 (Evidence application)
 評估結果 (Outcome evaluation)

提出問題 (Question formulation)

- **Therapy/Prevention** :
研究治療或預防方法的有效性
例如：服用“阿斯匹林”是否可以預防中風？
- **Harm/Etiology** :
研究暴露的危害或疾病的原因
例如：停經婦女使用荷爾蒙治療是否會增加乳癌的機曾？

EBM五步驟

EBM五步驟：
 提出問題 (Question formulation)
 搜尋證據 (Evidence search)
 嚴格判讀 (Critical appraisal)
 恰當運用 (Evidence application)
 評估結果 (Outcome evaluation)

提出問題 (Question formulation)

- **Diagnosis (tests)** :
研究檢查方法或臨床表徵對疾病診斷的有效性
例如：Low dose CT 診斷肺癌的敏感度及特異度為何？
- **Prognosis** :
建立疾病預後的預測模式，例如：利用 Ranson's criteria 預測急性胰臟炎死亡率為何？

EBM五步驟

EBM五步驟：
 提出問題 (Question formulation)
 搜尋證據 (Evidence search)
 嚴格判讀 (Critical appraisal)
 恰當運用 (Evidence application)
 評估結果 (Outcome evaluation)

提出問題 (Question formulation)

Patient	Intervention	Comparison	Outcome
---------	--------------	------------	---------

- P : patient and/or problem(病患)
- I : intervention(處理)
- C : comparison of intervention(對照)
- O : clinical outcome(臨床結果)

EBM五步驟

EBM五步驟：
 提出問題 (Question formulation)
 搜尋證據 (Evidence search)
 嚴格判讀 (Critical appraisal)
 恰當運用 (Evidence application)
 評估結果 (Outcome evaluation)

提出問題 (Question formulation)

Patient or Problem	Intervention	Comparison intervention	Outcome
Description of the patient or the target disorder of interest	Could include: -Exposure -Diagnostic test -Prognostic factor -Therapy -Patient perception etc.	Relevant most often when looking at therapy questions	Clinical outcome of interest to you and your patient

EBM五步驟

EBM五步驟：
 提出問題 (Question formulation)
 搜尋證據 (Evidence search)
 嚴格判讀 (Critical appraisal)
 恰當運用 (Evidence application)
 評估結果 (Outcome evaluation)

提出問題 (Question formulation)

Patient or Problem	Intervention	Comparison intervention	Outcome
65 year old man with stroke and moderate carotid stenosis	Carotid endarterectomy	Medical therapy	Functional Class

- In a 65 year old man with stroke and moderate carotid stenosis, can carotid endarterectomy improve functional class after stroke compared with medical therapy?

EBM五步驟

EBM五步驟：
 提出問題 (Question formulation)
 搜尋證據 (Evidence search)
 嚴格判讀 (Critical appraisal)
 恰當運用 (Evidence application)
 評估結果 (Outcome evaluation)

提出問題 (Question formulation)

陳先生因為有冠狀動脈家族病史，並且最近胸口悶悶的，因此要做相關檢查，經由自己查詢資料後得知可以經由心臟電腦斷層或心導管檢查，希望徵詢你的建議。

什麼是EBM?

請問你要用什麼關鍵字來搜尋?

- (1) 豬血糕, 熱狗
- (2) 冠狀動脈疾病, 心臟電腦斷層, 心導管
- (3) 胸口悶, 心臟電腦斷層
- (4) 大腸包小腸, 魯肉飯

EBM五步驟

EBM五步驟：
 提出問題 (Question formulation)
 搜尋證據 (Evidence search)
 嚴格判讀 (Critical appraisal)
 恰當運用 (Evidence application)
 評估結果 (Outcome evaluation)

● 提出問題 (Question formulation)

Patient	Intervention	Comparison	Outcome
病人	DSCT	心導管	診斷準確率

- P : patient and/or problem(病患)
- I : intervention(處理)
- C : comparison of intervention(對照)
- O : clinical outcome(臨床結果)

EBM五步驟

EBM五步驟：
 提出問題 (Question formulation)
 搜尋證據 (Evidence search)
 嚴格判讀 (Critical appraisal)
 恰當運用 (Evidence application)
 評估結果 (Outcome evaluation)

● 搜尋證據 (Evidence search)

- 關鍵字設定
- 資料庫選擇
- 學習各種資料庫的搜尋方法
- 多練習, 熟能生巧

EBM五步驟

EBM五步驟：
 提出問題 (Question formulation)
 搜尋證據 (Evidence search)
 嚴格判讀 (Critical appraisal)
 恰當運用 (Evidence application)
 評估結果 (Outcome evaluation)

● 搜尋證據 (Evidence search)

Primary Term	Synonym 1	Synonym 2	
P (OR	OR) AND
I (OR	OR) AND
C (OR	OR) AND
O (OR	OR)

- 關鍵字設定
- 資料庫選擇
- 學習各種資料庫的搜尋方法
- 多練習, 熟能生巧

EBM五步驟

EBM五步驟：
 提出問題 (Question formulation)
 搜尋證據 (Evidence search)
 嚴格判讀 (Critical appraisal)
 恰當運用 (Evidence application)
 評估結果 (Outcome evaluation)

● 搜尋證據 (Evidence search)

Category	Articles per day
Biomedical	5,000?
MEDLINE	1,500
Trials	95
Diagnostic?	-

- 關鍵字設定
- 資料庫選擇
- 學習各種資料庫的搜尋方法
- 多練習, 熟能生巧

EBM五步驟

EBM五步驟：
 提出問題 (Question formulation)


Secondary pre-appraised research

- Systems** • Computerised decision support
- Summaries** • Critically appraised topics, evidence-based guidelines
- Synopses** • Critically-appraised journal articles
- Syntheses** • Systematic reviews and meta-analysis

Primary original research

- Studies** • RCTs, Cohort studies, case-control studies, case series/reports

- 多練習, 熟能生巧



什麼是EBM?

- 請問你會從哪一個層級開始搜尋?

- (1) Studies
- (2) Syntheses
- (3) Synopses
- (4) Summaries

EBM五步驟

- 提出問題 (Question formulation)
- 搜尋證據 (Evidence search)
- 嚴格判讀 (Critical appraisal)
- 恰當運用 (Evidence application)
- 評估結果 (Outcome evaluation)

搜尋證據 (Evidence search)

UpToDate ONLINE 18.0

Search Results for "Intracranial aneurysm"

- All search results | Prioritize adult topics | Prioritize pediatric topics | Prioritize patient topics
- Unruptured intracranial aneurysms
- Screening for intracranial aneurysm
- Etiology, clinical manifestations, and diagnosis of aneurysmal subarachnoid hemorrhage
- Anticoagulant and antiplatelet therapy in patients with an unruptured intracranial aneurysm
- Treatment of aneurysmal subarachnoid hemorrhage**
 - Headache syndromes other than migraine
 - Extracranial manifestations of autosomal dominant polycystic kidney disease
 - Third cranial nerve (oculomotor nerve) palsy in adults
 - Spontaneous intracerebral hemorrhage: pathogenesis, clinical features, and diagnosis
 - Thunderclap headache
 - Recombinant human activated protein C in severe sepsis and septic shock
 - Nonischemic neurologic complications of congenital heart disease in adults
 - Carotid endarterectomy: Preoperative evaluation; surgical technique; and complications
 - Perimesencephalic nonaneurysmal subarachnoid hemorrhage

多練習，熟能生巧

EBM五步驟

- 提出問題 (Question formulation)
- 搜尋證據 (Evidence search)
- 嚴格判讀 (Critical appraisal)
- 恰當運用 (Evidence application)
- 評估結果 (Outcome evaluation)

搜尋證據 (Evidence search)

UpToDate ONLINE 18.0

Search Results for "Intracranial aneurysm"

Treatment of aneurysmal subarachnoid hemorrhage

DESCRIPTION - After aneurysmal SAH, the patient is at substantial risk of rebleeding: 3 to 4 percent in the first 24 hours and 1 to 2 percent each day in the first month (L1). Reoperation is associated with a mortality that is estimated to be 30 percent. Aneurysm repair is the only effective treatment to prevent rebleeding. (See Self, rebleeding, below.)

RELEVANCE - Surgery has been the mainstay of therapy of intracranial aneurysms; however, minimally-invasive techniques are becoming more widely used. In the future, gene therapy combined with endovascular techniques may offer improved results for aneurysm treatment (L2).

RECOMMENDATIONS - Surgical management of cerebral aneurysms is an effective and safe procedure with the evolution of microsurgical techniques in the hands of an experienced surgeon. Placement of a clip across the neck of the aneurysm remains the treatment of choice for most aneurysms (L1c, L2).

TEMPORARY VESSEL OCCLUSION - Temporary vessel occlusion using endovascular techniques such as hypothermia, induced hypertension, and temporary balloon catheter occlusion near the aneurysm (L2). Temporary clip occlusion may be achieved via traditional extracranial clip placement or with an intracranial balloon. Intraoperative angiography in a second diagnostic, subject to micro-Doppler angiography, is preferred to postoperative angiography after clip occlusion.

ENDOVASCULAR THERAPY - Endovascular therapy is a potential alternative to microsurgical clipping of intracranial aneurysms. In a pooled analysis of immediate endovascular placement, aneurysm clipping (of intracranial pressure was less than or equal to 30 mmHg without an neurological deficit or less than or equal to 80 mmHg in the presence of a skull) and aggressive postoperative hypertension, hypervolemia, and hemodilution therapy was evaluated in 35 patients with poor-grade aneurysmal SAH (L2). The outcome was reported as good in 16 patients (24 percent), fair in 11 (31 percent), and poor in 11 (31 percent); eight patients (23 percent) died. In contrast, 12 poor-grade aneurysms were treated with microsurgical clipping. Endovascular therapy was associated with a higher rate of rebleeding after vessel occlusion (L2).

RECOMMENDATIONS - Preoperative evaluation; surgical technique; and complications

- Perimesencephalic nonaneurysmal subarachnoid hemorrhage

多練習，熟能生巧

EBM五步驟

- 提出問題 (Question formulation)
- 搜尋證據 (Evidence search)
- 嚴格判讀 (Critical appraisal)
- 恰當運用 (Evidence application)
- 評估結果 (Outcome evaluation)

搜尋證據 (Evidence search)

ACP Journal Club - Search Results

Search for: intracranial aneurysm, clipping, coiling

Article type: Therapeutics (Diagnosis, Clinical Prediction Guide, Prognosis)

Search Help

Improve your results

The following words first appear in ACP Journal Club clipping, aneurysm. Rephrasing your query using different words may improve results.

Found 1 matches: Showing 1 - 1.

1. OAM - International subarachnoid aneurysm trial (ISAT) of neurosurgical clipping versus endovascular coiling in 2143 patients with ruptured intracranial aneurysms: a randomised comparison of effects on survival, dependency, seizures, rebleeding, subgroup, and aneurysm occlusion.

關鍵字設定
資料庫選擇
各種資料庫的搜尋方法
練習，熟能生巧

EBM五步驟

- 提出問題 (Question formulation)
- 搜尋證據 (Evidence search)
- 嚴格判讀 (Critical appraisal)
- 恰當運用 (Evidence application)
- 評估結果 (Outcome evaluation)

搜尋證據 (Evidence search)

THE COCHRANE LIBRARY

Independent high-quality evidence for health care decision making

Search Results

Found 6 records of RCTs available for "intracranial aneurysm and clipping and coiling, from 2003 to 2011 in Cochrane Database of Systematic Reviews"

1. OAM - International subarachnoid aneurysm trial (ISAT) of neurosurgical clipping versus endovascular coiling in 2143 patients with ruptured intracranial aneurysms: a randomised comparison of effects on survival, dependency, seizures, rebleeding, subgroup, and aneurysm occlusion.

關鍵字設定
資料庫選擇
學習各種資料庫的搜尋方法
多練習，熟能生巧

EBM五步驟

- 提出問題 (Question formulation)
- 搜尋證據 (Evidence search)
- 嚴格判讀 (Critical appraisal)
- 恰當運用 (Evidence application)
- 評估結果 (Outcome evaluation)

● 搜尋證據 (Evidence search)

EBM五步驟：

- 關鍵字設定
- 資料庫選擇
- 學習各種資料庫的搜尋方法
- 多練習，熟能生巧

EBM五步驟

- 提出問題 (Question formulation)
- 搜尋證據 (Evidence search)
- 嚴格判讀 (Critical appraisal)
- 恰當運用 (Evidence application)
- 評估結果 (Outcome evaluation)

● 搜尋證據 (Evidence search)

Try our newly improved Advanced Search

SEARCH PICO ADVANCED SEARCH

Population: patient
Intervention: TAE
Comparison: surgery
Outcome: outcome

EBM五步驟：

- 關鍵字設定
- 資料庫選擇
- 學習各種資料庫的搜尋方法
- 多練習，熟能生巧

EBM五步驟

- 提出問題 (Question formulation)
- 搜尋證據 (Evidence search)
- 嚴格判讀 (Critical appraisal)
- 恰當運用 (Evidence application)
- 評估結果 (Outcome evaluation)

● 搜尋證據 (Evidence search)

EBM五步驟：

- 關鍵字設定
- 資料庫選擇
- 學習各種資料庫的搜尋方法
- 多練習，熟能生巧

EBM五步驟

- 提出問題 (Question formulation)
- 搜尋證據 (Evidence search)
- 嚴格判讀 (Critical appraisal)
- 恰當運用 (Evidence application)
- 評估結果 (Outcome evaluation)

● 搜尋證據 (Evidence search)

EBM五步驟：

- 關鍵字設定
- 資料庫選擇
- 學習各種資料庫的搜尋方法
- 多練習，熟能生巧

EBM五步驟

- 提出問題 (Question formulation)
- 搜尋證據 (Evidence search)
- 嚴格判讀 (Critical appraisal)
- 恰當運用 (Evidence application)
- 評估結果 (Outcome evaluation)

● 搜尋證據 (Evidence search)

EBM五步驟：

- 關鍵字設定
- 資料庫選擇
- 學習各種資料庫的搜尋方法
- 多練習，熟能生巧

EBM五步驟

- 提出問題 (Question formulation)
- 搜尋證據 (Evidence search)
- 嚴格判讀 (Critical appraisal)
- 恰當運用 (Evidence application)
- 評估結果 (Outcome evaluation)

● 搜尋證據 (Evidence search)

METHODS

Data sources: Medline, PsycInfo, Cochrane Central Register of Controlled Trials, and Cochrane Database of Systematic Reviews to March 2013; clinical trial registries; reference lists; and technical advisors.

Study Selection: We examined controlled clinical trials and observational studies comparing screening to no screening, and controlled clinical trials comparing different screening intervals. We also examined controlled clinical trials and observational studies comparing one of the following active treatments to conservative treatment in patients with early-stage HCC: transarterial chemoembolization (TACE), partial hepatic resection, orthotopic liver transplant (OLT), radiofrequency ablation (RFA), and sorafenib. Because of the dearth of studies for all treatments other than TACE comparing active to conservative treatments, we evaluated noncomparative observational studies for evidence on harms and long-term survival. The population of interest was patients with chronic liver disease with or without cirrhosis.

EBM五步驟：

- 關鍵字設定
- 資料庫選擇
- 學習各種資料庫的搜尋方法
- 多練習，熟能生巧

EBM五步驟：

- 提出問題 (Question formulation)
- 搜尋證據 (Evidence search)
- 嚴格判讀 (Critical appraisal)
- 恰當運用 (Evidence application)
- 評估結果 (Outcome evaluation)

Effects of screening on mortality: RCTs

Two trials, both conducted in China, compared the effects of screening to no screening on mortality among participants mainly with hepatitis B. One trial used a cluster-randomized design to assign factories, business, and schools to screening or no screening groups. Screening group participants (n = 9,757) were offered serum AFP testing and ultrasonography every 6 months. The primary outcome of HCC mortality occurred less frequently in the screening group (83.2/100,000 person-years vs 131.5/100,000 person-years; rate ratio 0.63, 95% CI 0.41–0.98). However, this trial, carried a high-risk of bias because of several serious methodological limitations that threaten the validity of the results. The second trial used patient-level randomization stratified by township to assign hepatitis B patients to the screening intervention (n = 3,712), which consisted of serial AFP tests followed by ultrasound for high AFP values, or the usual care group (n = 1,869). HCC mortality was similar in both groups (1,138/100,000 person-years vs 1114/100,000 person-years; p = 0.86), as was all-cause mortality (1,843/100,000 person-years vs 1,788/100,000 person-years, p = NS). This trial carried an unclear risk of bias because of poor reporting of randomization and allocation concealment techniques.

Two additional trials compared different ultrasound screening intervals. One unclear risk of bias trial found no survival advantage comparing 4-month to 12-month ultrasound screening intervals in patients with serologic evidence of hepatitis B or C. A trial with low risk-of-bias compared 3-month to 6-month ultrasound screening intervals in 1,278 patients with cirrhosis from alcohol use and/or viral hepatitis and found similar all-cause mortality rates in both groups.

- 關鍵字設定
- 資料庫選擇
- 學習各種資料庫的搜尋方法
- 多練習，熟能生巧

EBM五步驟：

- 提出問題 (Question formulation)
- 搜尋證據 (Evidence search)
- 嚴格判讀 (Critical appraisal)
- 恰當運用 (Evidence application)
- 評估結果 (Outcome evaluation)

EBM五步驟

Oxford Centre for Evidence-Based Medicine 2011 Levels of Evidence

Question	Step 1 (Level 1*)	Step 2 (Level 2*)	Step 3 (Level 3*)	Step 4 (Level 4*)	Step 5 (Level 5)
How common is the problem?	Local and current random sample surveys (or censuses)	Systematic review of surveys that allow matching to local circumstances**	Local non-random sample**	Case-series**	n/a
Is this diagnostic or monitoring test accurate? (Diagnosis)	Systematic review of cross-sectional studies with consistently applied reference standard and blinding	Individual cross-sectional studies with consistently applied reference standard and blinding	Non-consecutive studies, or studies without consistently applied reference standards**	Case-control studies, or poor or non-independent reference standards**	Mechanism-based reasoning
What will happen if we do not add a therapy? (Prognosis)	Systematic review of inception cohort studies	Inception cohort studies	Cohort study or control arm of randomized trial**	Case-series or case-control studies, or poor quality prognostic cohort study**	n/a
Does this intervention help? (Treatment Benefits)	Systematic review of randomized trials or n-of-1 trials	Randomized trial or observational study with dramatic effect	Non-randomized controlled cohort/follow-up study**	Case-series, case-control studies, or historically controlled studies**	Mechanism-based reasoning
What are the COMMON harms? (Treatment Harms)	Systematic review of randomized trials, systematic review of nested case-control studies, n-of-1 trials with the patient you are raising the question about, or observational study with dramatic effect	Individual randomized trial or (exceptionally) observational study with dramatic effect	Non-randomized controlled cohort/follow-up study (post-marketing surveillance) provided there are sufficient numbers to rule out a common harm. (For long-term harms the duration of follow-up must be sufficient.)**	Case-series, case-control, or historically controlled studies**	Mechanism-based reasoning
What are the RARE harms? (Treatment Harms)	Systematic review of randomized trials or n-of-1 trial	Randomized trial or (exceptionally) observational study with dramatic effect	Non-randomized controlled cohort/follow-up study**	Case-series, case-control, or historically controlled studies**	Mechanism-based reasoning
Is this (early detection) test worthwhile? (Screening)	Systematic review of randomized trials	Randomized trial	Non-randomized controlled cohort/follow-up study**	Case-series, case-control, or historically controlled studies**	Mechanism-based reasoning

- 關鍵字設定
- 資料庫選擇
- 學習各種資料庫的搜尋方法
- 多練習，熟能生巧

Expert

EBM五步驟：

- 提出問題 (Question formulation)
- 搜尋證據 (Evidence search)
- 嚴格判讀 (Critical appraisal)
- 恰當運用 (Evidence application)
- 評估結果 (Outcome evaluation)

EBM五步驟

Oxford Centre for Evidence-Based Medicine 2011 Levels of Evidence

Question	Step 1 (Level 1*)	Step 2 (Level 2*)	Step 3 (Level 3*)	Step 4 (Level 4*)	Step 5 (Level 5)
How common is the problem?	Local and current random sample surveys (or censuses)	Systematic review of surveys that allow matching to local circumstances**	Local non-random sample**	Case-series**	n/a
Is this diagnostic or monitoring test accurate? (Diagnosis)	Systematic review of cross-sectional studies with consistently applied reference standard and blinding	Individual cross-sectional studies, or studies without consistently applied reference standards**	Case-control studies, or poor or non-independent reference standards**	Case-series or case-control studies, or poor quality prognostic cohort study**	Mechanism-based reasoning
What will happen if we do not add a therapy? (Prognosis)	Systematic review of inception cohort studies	Inception cohort studies	Cohort study or control arm of randomized trial**	Case-series or case-control studies, or poor quality prognostic cohort study**	n/a
Does this intervention help? (Treatment Benefits)	Systematic review of randomized trials or n-of-1 trials	Randomized trial or observational study with dramatic effect	Non-randomized controlled cohort/follow-up study**	Case-series, case-control studies, or historically controlled studies**	Mechanism-based reasoning
What are the COMMON harms? (Treatment Harms)	Systematic review of randomized trials, systematic review of nested case-control studies, n-of-1 trials with the patient you are raising the question about, or observational study with dramatic effect	Individual randomized trial or (exceptionally) observational study with dramatic effect	Non-randomized controlled cohort/follow-up study (post-marketing surveillance) provided there are sufficient numbers to rule out a common harm. (For long-term harms the duration of follow-up must be sufficient.)**	Case-series, case-control, or historically controlled studies**	Mechanism-based reasoning
What are the RARE harms? (Treatment Harms)	Systematic review of randomized trials or n-of-1 trial	Randomized trial or (exceptionally) observational study with dramatic effect	Non-randomized controlled cohort/follow-up study**	Case-series, case-control, or historically controlled studies**	Mechanism-based reasoning
Is this (early detection) test worthwhile? (Screening)	Systematic review of randomized trials	Randomized trial	Non-randomized controlled cohort/follow-up study**	Case-series, case-control, or historically controlled studies**	Mechanism-based reasoning


Systematic review (highlighted in pink)

Cohort studies (highlighted in yellow)

Randomized (highlighted in green)

Case-control study (highlighted in blue)

- 關鍵字設定
- 資料庫選擇
- 學習各種資料庫的搜尋方法
- 多練習，熟能生巧



EBM五步驟：

- 提出問題 (Question formulation)
- 搜尋證據 (Evidence search)
- 嚴格判讀 (Critical appraisal)
- 恰當運用 (Evidence application)
- 評估結果 (Outcome evaluation)

EBM五步驟

longitudinal studies - Cohort studies

Prospective

Retrospective

2,000 2,001 2,002 2,003 2,004 2,005 2,006 2,007

cross-sectional studies

Patient → Random

→ Not Random

- 關鍵字設定
- 資料庫選擇
- 學習各種資料庫的搜尋方法
- 多練習，熟能生巧

EBM五步驟：

- 提出問題 (Question formulation)
- 搜尋證據 (Evidence search)
- 嚴格判讀 (Critical appraisal)
- 恰當運用 (Evidence application)
- 評估結果 (Outcome evaluation)

research design

PubMed.gov

Search: Published

Limits: Advanced search Help

Early Hum Dev. 2011 Mar 10. [Epub ahead of print]

Longitudinal study of smoking cessation before pregnancy and children's cognitive abilities at 60 months of age.

Heinonen K, Räikkönen K, Pesonen AK, Andersson S, Kajantie E, Eriksson JG, Wolkie D, Lano A. Institute of Behavioral Science, P.O. Box 9, FI-00014, University of Helsinki, Finland.

Abstract

BACKGROUND: An inverse relationship exists between the rates of maternal smoking during pregnancy and children's cognitive abilities. The effect of maternal cessation of smoking before pregnancy on child's cognitive development is less clear.

AIMS: To study whether maternal cessation of smoking before pregnancy is associated with children's cognitive abilities.

STUDY DESIGN AND SUBJECTS: The original cohort included all 1535 live-born infants admitted to the neonatal wards during 1 year and 858 randomly recruited non-admitted infants. The present study sample comprised 1019 (66.2%) children of the original sample born at term and free of any major impairment followed up to 60 months.

OUTCOME MEASURES: Child's general reasoning, visual-motor integration, verbal competence, and language comprehension

- 關鍵字設定
- 資料庫選擇
- 學習各種資料庫的搜尋方法
- 多練習，熟能生巧

EBM五步驟：
 提出問題 (Question formulation)
 搜尋證據 (Evidence search)
 嚴格判讀 (Critical appraisal)
 恰當運用 (Evidence application)
 評估結果 (Outcome evaluation)

research design

PubMed.gov
 Search: PubMed Limits Advanced search Help

Display Settings Abstract Send to

Eur J Cardiovasc Nurs, 2011 Mar 11. [Epub ahead of print]

The impact of the metabolic syndrome on health-related quality of life: A cross-sectional study in Greece.

Tziallas D, Kastanioti C, Kostapanos MS, Skapinakis P, Elisaf MS, Mavreas V.
 Department of Internal Medicine, University of Ioannina, Ioannina, Greece.

Abstract
BACKGROUND: Metabolic syndrome (MetS) is a chronic, progressive and multi-complex health problem that can trigger physical, emotional and psychosocial problems. The aim of this study is to investigate the association between MetS and health-related quality of life (HRQoL) as well as depressive and anxiety disorders.
METHODS: New consecutive patients who attended an outpatient lipid clinic for evaluation for MetS were eligible for inclusion in the study. The MetS was defined according to the new definition of International Diabetes Federation (IDF). The Medical Outcomes Study, Short-Form-36 (SF-36) was used to assess HRQoL. Anxiety and depressive symptoms were assessed by a validated Greek version of the Hospital Anxiety and Depression Scale (HADS).

多練習 · 熟能生巧

EBM五步驟：
 提出問題 (Question formulation)
 搜尋證據 (Evidence search)
 嚴格判讀 (Critical appraisal)
 恰當運用 (Evidence application)
 評估結果 (Outcome evaluation)

research design

PubMed.gov
 Search: PubMed Limits Advanced search Help

Display Settings Abstract Send to

Osteoarthritis Cartilage, 2011 Mar 8. [Epub ahead of print]

Systematic review of the concurrent and predictive validity of MRI biomarkers in OA.

Hunter DJ, Zhang W, Conaghan PG, Hirkko K, Menashe L, Li L, Reichmann WM, Losina E.
 Rheumatology Department, Royal North Shore Hospital and Northern Clinical School, University of Sydney, Sydney, NSW Australia; Division of Research, New England Baptist Hospital, Boston MA, USA.

Abstract
OBJECTIVE: To summarize literature on the concurrent and predictive validity of MRI-based measures of osteoarthritis (OA) structural change.
METHODS: An online literature search was conducted of the OVID, EMBASE, CINAHL, PsycInfo and Cochrane databases of articles published up to the time of the search, April 2009. 1,338 abstracts obtained with this search were preliminarily screened for relevance by two reviewers. Of these, 243 were selected for data extraction for this analysis on validity as well as separate reviews on discriminative validity and diagnostic performance. Of these 142 manuscripts included data pertinent to concurrent validity, and 81 manuscripts for the predictive validity review. For this analysis we extracted data on criterion (concurrent and predictive) validity from both longitudinal and cross-sectional studies for all synovial joint tissues as it relates to MRI

多練習 · 熟能生巧


● 搜尋證據 (Evidence search)
 原始研究 (Original Study)

■ 隨機對照實驗 (Randomized Control Trial, RCT)

給實驗組及對照組不同的治療，觀察其後果。
 例如：實驗組吃aspirin，對照組吃澱粉，比較兩組五年後中風的機率。為目前臨床流行病學中公認證據力最強之原始介入性研究設計 (primary interventional study)。在此設計中，如果能夠採行雙盲對照 (double blinded, placebo-controlled) 則干擾因素的影響可以進一步減少

```

    graph TD
      A[民眾] --> B[Randomized]
      B --> C[實驗組]
      B --> D[對照組]
      C --> E[aspirin]
      D --> F[澱粉]
      E --> G[五年後中風機率?]
      F --> G
    
```




● 搜尋證據 (Evidence search)
 原始研究 (Original Study)

■ 世代研究 (Cohort Study)

觀察自然暴露/治療方式的影響，長期追蹤其結果。例如：比較金山鄉有吃aspirin及沒有吃aspirin的人，五年後新發中風的機率。提供比隨機對照試驗證據力較次一級之研究證據，是屬於臨床流行病學中之觀察性研究。針對幾個子群，由接受暴露因子開始，一直追蹤到結果。

```

    graph TD
      A[金山鄉] --> B[吃 aspirin]
      A --> C[沒吃 aspirin]
      B --> D[五年後中風機率?]
      C --> D
    
```



● 搜尋證據 (Evidence search)

原始研究 (Original Study)

■ 病例对照研究 (Case Control Study)

選擇病人組及對照組，研究其暴露/治療的影響。
例如：比較500個有中風的個案及500個沒有中風的個案，他們吃 aspirin 的比例。這是**流行病學中極重要的研究方式**，雖然它的臨床**證據力**較世代研究為低，但可以在**短時間內**以**少許資金與努力**，就能得到重要的科學發現，是最有效率的研究設計，但也容易受到偏見的影響。



● 搜尋證據 (Evidence search)

原始研究 (Original Study)

■ 個案報告 (Case Report)

收集**同一臨床特色之病人**，回顧文獻報告加以比較。
例如：收集5位吃aspirin預防中風結果發生**消化道出血**的病人報告及其臨床特徵，並回顧相關文獻，缺點是**缺乏對照組**，且**證據力也較低**。



● 搜尋證據 (Evidence search)

文獻回顧 (Review of Literature)

■ 系統性回顧 (systemic review)

系統性的文獻回顧，將個人好惡及偏差減至最低。**需引用各種文獻資料庫**，並說明查詢的關鍵字，**有引用文獻的明確標準**，不能依作者喜好選取文獻，通常會使用統計方法 (**Meta-Analysis 統合分析**)，以解決臨床爭議為目的。

● 搜尋證據 (Evidence search)

文獻回顧 (Review of Literature)

■ 非系統性回顧 (non-systemic review)

傳統文獻回顧，**容易有個人好惡及偏差**。以作者的判斷引用文獻，通常沒有使用特殊的統計方法，以教育臨床醫師為目的。

EBM五步驟

- 提出問題 (Question formulation)
- 搜尋證據 (Evidence search)
- 嚴格判讀 (Critical appraisal)
- 恰當運用 (Evidence application)
- 評估結果 (Outcome evaluation)

● 搜尋證據 (Evidence search)



- 關鍵字設定
- 資料庫選擇
- 學習各種資料庫的搜尋方法
- 多練習，熟能生巧

EBM五步驟

- 提出問題 (Question formulation)
- 搜尋證據 (Evidence search)
- 嚴格判讀 (Critical appraisal)
- 恰當運用 (Evidence application)
- 評估結果 (Outcome evaluation)

● 搜尋證據 (Evidence search)

■ 請問你個人比較喜歡的可樂是？

- (1) 可口可樂
- (2) 百事可樂
- (3) 他牌可樂

- 關鍵字設定
- 資料庫選擇
- 學習各種資料庫的搜尋方法
- 多練習，熟能生巧

EBM五步驟

- 提出問題 (Question formulation)
- 搜尋證據 (Evidence search)
- 嚴格判讀 (Critical appraisal)
- 恰當運用 (Evidence application)
- 評估結果 (Outcome evaluation)

● 搜尋證據 (Evidence search)



- 關鍵字設定
- 資料庫選擇
- 學習各種資料庫的搜尋方法
- 多練習，熟能生巧

EBM五步驟

- 提出問題 (Question formulation)
- 搜尋證據 (Evidence search)
- 嚴格判讀 (Critical appraisal)
- 恰當運用 (Evidence application)
- 評估結果 (Outcome evaluation)

● 搜尋證據 (Evidence search)



- 關鍵字設定
- 資料庫選擇
- 學習各種資料庫的搜尋方法
- 多練習，熟能生巧

EBM五步驟

- 提出問題 (Question formulation)
- 搜尋證據 (Evidence search)
- 嚴格判讀 (Critical appraisal)
- 恰當運用 (Evidence application)
- 評估結果 (Outcome evaluation)

● 搜尋證據 (Evidence search)

■ 請問研究結果中試驗者比較喜歡的可樂是？

- (1) 可口可樂
- (2) 百事可樂
- (3) 他牌可樂

- 關鍵字設定
- 資料庫選擇
- 學習各種資料庫的搜尋方法
- 多練習，熟能生巧

EBM五步驟

- 提出問題 (Question formulation)
- 搜尋證據 (Evidence search)
- 嚴格判讀 (Critical appraisal)
- 恰當運用 (Evidence application)
- 評估結果 (Outcome evaluation)

● 搜尋證據 (Evidence search)



- 關鍵字設定
- 資料庫選擇
- 學習各種資料庫的搜尋方法
- 多練習，熟能生巧

EBM五步驟

EBM五步驟：
 提出問題 (Question formulation)
 搜尋證據 (Evidence search)
 嚴格判讀 (Critical appraisal)
 恰當運用 (Evidence application)
 評估結果 (Outcome evaluation)

- 搜尋證據 (Evidence search)
- 請問研究結果中試驗者比較喜歡的可樂是？
 - (1) 可口可樂
 - (2) 百事可樂
 - (3) 他牌可樂

■ 關鍵字設定
 ■ 資料庫選擇
 ■ 學習各種資料庫的搜尋方法
 ■ 多練習，熟能生巧

EBM五步驟

EBM五步驟：
 提出問題 (Question formulation)
 搜尋證據 (Evidence search)
 嚴格判讀 (Critical appraisal)
 恰當運用 (Evidence application)
 評估結果 (Outcome evaluation)

- 嚴格判讀 (Critical appraisal)-簡單判讀
- **Oxford Evidence-Based Medicine Levels-Level 1-5**

最主要有三個主要步驟，即為**VIP**：

- **V** (Validity/Reliability) 效度/信度：文獻研究收集個案方式是否**嚴謹**，是否具有**有效度(Validity)**？
- **I** (Importance/Impact) 重要性、文獻統計方法是**否正確**，**結果是否具有重要性**？
- **P** (Practice/Applicability) 臨床適用性。

EBM五步驟

EBM五步驟：
 提出問題 (Question formulation)
 搜尋證據 (Evidence search)
 嚴格判讀 (Critical appraisal)
 恰當運用 (Evidence application)
 評估結果 (Outcome evaluation)

V

- 嚴格判讀 (Critical appraisal)
- 文獻個案收集方式是否**隨機取樣** (Randomization)?
Randomization、case-control study、case-series
- 個案**追蹤成功率**是否高？
- 是否進行**單盲**(Single-blind)或**雙盲**(Double-blind)測試？
- 研究進行中個案是否經由**嚴謹管制**？
- 個案是否**追蹤遺失**？
 - 文獻研究收集個案方式是否嚴謹，是否具有有效度(Validity)?
 - 文獻統計方法是否正確，結果是否具有重要性(Importance)?
 - 文獻是否適合應用在您的病人身上?

EBM五步驟

EBM五步驟：
 提出問題 (Question formulation)
 搜尋證據 (Evidence search)
 嚴格判讀 (Critical appraisal)
 恰當運用 (Evidence application)
 評估結果 (Outcome evaluation)

I

不同的問題利用不同的統計量

Therapy/Prevention	Diagnosis
RRR	Sensitivity
ARR	Specificity
NNT	PPV
Relative risk	NPV
Odds ratio(Case-control study)	LR

- 文獻研究收集個案方式是否嚴謹，是否具有有效度(Validity)?
- 文獻統計方法是否正確，結果是否具有重要性(Importance)?
- 文獻是否適合應用在您的病人身上?

EBM五步驟

EBM五步驟：
 提出問題 (Question formulation)
 搜尋證據 (Evidence search)
 嚴格判讀 (Critical appraisal)
 恰當運用 (Evidence application)
 評估結果 (Outcome evaluation)

I

Diagnostic test	Disease	
	Present(有病)	Absent(沒病)
Positive	731 <small>a</small>	270 <small>b</small>
Negative	78 <small>c</small>	1500 <small>d</small>

Sensitivity (敏感度) 為**有病者**診斷結果為**陽性**的比率=真陽性率=真陽性 / 生病 = $a / a+c = 731/809 = 90\%$

Specificity (特異性) 為**沒病者**診斷結果為**陰性**的比率
 $d/b+d = \text{真陰性率} = \text{真陰性} / \text{健康} = d / b+d = 1500/1770 = 85\%$

- 文獻統計方法是否正確，結果是否具有重要性(Importance)?
- 文獻是否適合應用在您的病人身上?

EBM五步驟

EBM五步驟：
 提出問題 (Question formulation)
 搜尋證據 (Evidence search)
 嚴格判讀 (Critical appraisal)
 恰當運用 (Evidence application)
 評估結果 (Outcome evaluation)

I

Diagnostic test	Disease	
	Present(有病)	Absent(沒病)
Positive	731 <small>a</small>	270 <small>b</small>
Negative	78 <small>c</small>	1500 <small>d</small>

Positive Predictive Value, PPV (陽性預測值) 診斷試驗結果呈現**陽性且確實有病者**的比率=真陽性 / 陽性試驗結果 = $a / a+b = 731/1001 = 73\%$

Negative Predictive Value, NPV (陰性預測值) 診斷試驗結果呈**陰性且確實無患病者**的比率=真陰性 / 陰性試驗結果 = $d / c+d = 1500/1578 = 95\%$

- 文獻研究收集個案方式是否嚴謹，是否具有有效度(Validity)?
- 文獻統計方法是否正確，結果是否具有重要性(Importance)?
- 文獻是否適合應用在您的病人身上?

I EBM五步驟

EBM五步驟：
 提出問題 (Question formulation)
 搜尋證據 (Evidence search)
 嚴格判讀 (Critical appraisal)
 恰當運用 (Evidence application)

Diagnostic test	Disease	
	Present(有病)	Absent(沒病)
Positive	73 ^a	270 ^b
Negative	78 ^c	1500 ^d

prevalence $a+c/a+b+c+d=31\%$

Likelihood Ratios (相似比)
 分子：疾病中診斷試驗(陽性或陰性)比率
 分母：無疾病中診斷試驗(陽性或陰性)比率

LR(+) **真陽性率 / 假陽性率**=Sensitivity / (1-Specificity) = $(a/a+c) / (b/b+d) = 90\% / 15\% = 6$

LR(-) **假陰性率 / 真陰性率**=(1-Sensitivity) / Specificity = $(c/a+c) / (d/b+d) = 10\% / 85\% = 0.12$

■ 文獻是否適合應用在您病人身上?

I EBM五步驟

EBM五步驟：
 提出問題 (Question formulation)
 搜尋證據 (Evidence search)
 嚴格判讀 (Critical appraisal)
 恰當運用 (Evidence application)

Treatment	Event	
	Positive	Negative
experimental -(RCT 或 Cohort studies)	A = 1	B = 29
control -(RCT 或 Cohort studies)	C = 9	D = 21

實驗組事件發生率 (EER, Experimental Event Rate) $a/a+b = 0.033$

對照組事件發生率 (CER, Control Event Rate) $c/c+d = 0.30$

相對風險比率差 (RRR, Relative Risk Reduction) $[EER - CER] / CER$ 接受治療組比未接受治療的對照組間，不良結果機率下降之比例，伴隨95%信賴區間(CI)

■ 文獻是否適合應用在您病人身上?

I EBM五步驟

EBM五步驟：
 提出問題 (Question formulation)
 搜尋證據 (Evidence search)
 嚴格判讀 (Critical appraisal)
 恰當運用 (Evidence application)

Treatment	Event	
	Positive	Negative
experimental -(RCT 或 Cohort studies)	A = 1	B = 29
control -(RCT 或 Cohort studies)	C = 9	D = 21

絕對風險比率差 (ARR, Absolute Risk Reduction) $EER - CER = |0.033 - 0.3| = 0.267$
 治療組與對照組間不良結果機率差的絕對值，伴隨95%信賴區間(CI)

需要被治療的病人數目 (NNT, Number Needed to Treat) $1/ARR = 1/0.267 = 3.75$ ，為減少一個不良結果所需治療4病人，伴隨95%信賴區間(CI)

■ 文獻研究收集個案方式是否嚴謹，是否具有效度(Validity)?
 ■ 文獻統計方法是否正確，結果是否具有重要性(Importance)?
 ■ 文獻是否適合應用在您病人身上?

I EBM五步驟

EBM五步驟：
 提出問題 (Question formulation)
 搜尋證據 (Evidence search)
 嚴格判讀 (Critical appraisal)
 恰當運用 (Evidence application)

Treatment	Event	
	Positive	Negative
experimental -(RCT 或 Cohort studies)	A = 1	B = 29
control -(RCT 或 Cohort studies)	C = 9	D = 21

Relative risk (RR) -相對危險比 $EER/CER = (a/a+b)/(c/c+d) = 0.11$ ，用於隨機對照試驗(RCT)與世代研究(cohort study)

■ 文獻研究收集個案方式是否嚴謹，是否具有效度(Validity)?
 ■ 文獻統計方法是否正確，結果是否具有重要性(Importance)?
 ■ 文獻是否適合應用在您病人身上?

I EBM五步驟

EBM五步驟：
 提出問題 (Question formulation)
 搜尋證據 (Evidence search)
 嚴格判讀 (Critical appraisal)
 恰當運用 (Evidence application)

Treatment	Event	
	Positive	Negative
Exposed (Case-control study)	A = 1	B = 29
Not exposed (Case-control study)	C = 9	D = 21

Experimental event Odds $a/b = 0.034$

Control event Odds $c/d = 0.43$

Odds ratio-勝算比 $(a/b)/(c/d) = ad/bc = 0.08$ ，通常被使用於個案對照研究之中。為試驗組中發生結果的勝算(Odds)與對照組中發生結果的勝算，此兩者間的比值就稱為勝算比(OR)

■ 文獻研究收集個案方式是否嚴謹，是否具有效度(Validity)?
 ■ 文獻統計方法是否正確，結果是否具有重要性(Importance)?
 ■ 文獻是否適合應用在您病人身上?

I EBM五步驟

EBM五步驟：
 提出問題 (Question formulation)
 搜尋證據 (Evidence search)
 嚴格判讀 (Critical appraisal)
 恰當運用 (Evidence application)
 評估結果 (Outcome evaluation)

- 嚴格判讀 (Critical appraisal)-有深度判讀-CASP
- Systematic reviews之評讀

Q1. Did the review ask a clearly-focused question?

Q2. Did the review include the right type of study?

Q3. Did the reviewers try to identify all relevant studies?

■ 文獻研究收集個案方式是否嚴謹，是否具有效度(Validity)?
 ■ 文獻統計方法是否正確，結果是否具有重要性(Importance)?
 ■ 文獻是否適合應用在您病人身上?

I **EBM五步驟**

EBM五步驟：
 提出問題 (Question formulation)
 搜尋證據 (Evidence search)
 嚴格判讀 (Critical appraisal)
 恰當運用 (Evidence application)
 評估結果 (Outcome evaluation)

- 嚴格判讀 (Critical appraisal)-有深度判讀
- Systematic reviews之評讀

Q4. Did the reviewers assess the quality of the included studies?

Q5. If the results of the studies have been combined, was it reasonable to do so?

Q6. How are the results presented and what is the main result?

Q7. How precise are these results?

- 文獻研究收集個案方式是否嚴謹，是否具有效度 (Validity)?
- 文獻統計方法是否正確，結果是否具有重要性 (Importance)?
- 文獻是否適合應用在你的人身上?

I **EBM五步驟**

EBM五步驟：
 提出問題 (Question formulation)
 搜尋證據 (Evidence search)
 嚴格判讀 (Critical appraisal)
 恰當運用 (Evidence application)
 評估結果 (Outcome evaluation)

- 嚴格判讀 (Critical appraisal)-有深度判讀
- Systematic reviews之評讀

Q8. Can the results be applied to the local population?

Q9. Were all important outcomes considered?

Q10. Should policy or practice change as a result of the evidence contained in this review?

- 文獻研究收集個案方式是否嚴謹，是否具有效度 (Validity)?
- 文獻統計方法是否正確，結果是否具有重要性 (Importance)?
- 文獻是否適合應用在你的人身上?

P **EBM五步驟**

EBM五步驟：
 提出問題 (Question formulation)
 搜尋證據 (Evidence search)
 嚴格判讀 (Critical appraisal)
 恰當運用 (Evidence application)
 評估結果 (Outcome evaluation)

- 嚴格判讀 (Critical appraisal)-有深度判讀
- P (Practice/Applicability) 臨床適用性。

如果我們相信這個研究，它的結果**是否可以應用**在我們的**病患身上**？”其中要考量的是病患的差異、可運用的資源以及病患的偏好。在這個部份我們可以考量病患的生物因素 (biologic issues) 即是”同樣的治療應用在**不同的病患族群**是否有不同的反應？”

- 文獻研究收集個案方式是否嚴謹，是否具有效度 (Validity)?
- 文獻統計方法是否正確，結果是否具有重要性 (Importance)?
- 文獻是否適合應用在你的人身上?

P **EBM五步驟**

EBM五步驟：
 提出問題 (Question formulation)
 搜尋證據 (Evidence search)
 嚴格判讀 (Critical appraisal)
 恰當運用 (Evidence application)
 評估結果 (Outcome evaluation)

- 恰當運用 (Evidence application)

Evidence is never enough
 證據，永遠是不足的

Can the results be applied to my patient?
 -有效且重要的證據**能實際應用**到我照顧的**病人**身上嗎？

Evidence Applicability -證據適用性

- 實際應用到你的病人
- 將證據“外推”，應用於個案？
- 個案自己的價值觀與喜好？

P **EBM五步驟**

EBM五步驟：
 提出問題 (Question formulation)
 搜尋證據 (Evidence search)
 嚴格判讀 (Critical appraisal)
 恰當運用 (Evidence application)
 評估結果 (Outcome evaluation)

- 恰當運用 (Evidence application)

病人與研究文獻所收錄的病人，有無明顯差異？
 - 研究族群為**60-70歲**有效，但是你的病人為**40歲**

評估病人接受與研究文獻所施行**相同的治療** (或處置)，所得**益處** (及害處) 將有多大？
 -應用到您的個案前，需先藉此**修正文獻結果**
 -單位或是你自己**是否有能力 (資源) 施行相同的治療** (或處置)？執行品質為何？

- 實際應用到你的病人
- 將證據“外推”，應用於個案？
- 個案自己的價值觀與喜好？

P **EBM五步驟**

EBM五步驟：
 提出問題 (Question formulation)
 搜尋證據 (Evidence search)
 嚴格判讀 (Critical appraisal)
 恰當運用 (Evidence application)
 評估結果 (Outcome evaluation)

- 恰當運用 (Evidence application)

是否清楚與個案 (及家屬) **說明溝通**？

個案對治療 (或處置) 的**遵從性**？

實際**追蹤**你的個案的**成果 (outcome)**

- 實際應用到你的病人
- 將證據“外推”，應用於個案？
- 個案自己的價值觀與喜好？

EBM五步驟

EBM五步驟：
 提出問題 (Question formulation)
 搜尋證據 (Evidence search)
 嚴格判讀 (Critical appraisal)
 恰當運用 (Evidence application)
 評估結果 (Outcome evaluation)

● **評估結果 (Outcome evaluation)**

■ **Self - Evaluation** (自我評估)

- 所遇到的臨床問題是**可以回答**的嗎？
- 搜尋文獻及評讀證據的**速度夠快**嗎？
- 這些證據應用在**適當的臨床病人**嗎

■ **Evaluation (Audit) by Expert or peer**

- **專家監督**或同儕彼此評估

參考文獻

1. 楊培銘：實證醫學與健康照護。臺灣醫學 2004; 8: 862-4。
2. 陳杰峰、蔡宛真、邱文達：實證醫學於健康照護之應用。臺灣醫學 2004; 8: 235-40。
3. Sackett DL, Rosenberg WM, Gray JA, et al: Evidence based medicine: what it is and what it isn't. BMJ 1996; 312: 71-2.
4. Levels of evidence. Oxford Centre for Evidence-Based Medicine. Web site.

參考文獻

5. 林愉珊：實證醫學相關資料庫檢索策略之探討。臺灣醫學2003; 7: 568-74。
2. 陳杰峰、蔡宛真、邱文達：實證醫學於健康照護之應用。臺灣醫學，2004; 8: 235-40。
6. Straus SE, Sackett DL: Using research findings in clinical practice. BMJ 1998; 317: 339-42.
7. Staunton M: Evidence-based radiology: steps 1 and 2-asking answerable questions and searching for evidence. Radiology, 2007; 242: 23-31.

參考文獻

8. Barratt A, Wyer PC, Hatala R, et al: Tips for learners of evidence-based medicine: Relative risk reduction, absolute risk reduction and number needed to treat. CMAJ. 2004; 171: 353-8.
9. Haynes RB: Of studies, syntheses, synopses, summaries, and systems: the "5S" evolution of information services for evidence-based healthcare decisions. Evid Based Med 2006; 11: 162-4.

Thank's for your attention